

After Cosmetic Reconstruction

It usually takes time to adjust to the feel of your new bite. When the bite is altered or the position of the teeth is changed it takes several days for the brain to recognize the new position of your teeth or their thickness as normal. If you continue to detect any high spots or problems with your bite, call us so we can schedule an adjustment appointment.

It is normal to experience some hot and cold sensitivity. The teeth require some time to heal after removal of tooth structure and will be sensitive in the interim. Your gums may also be sore for a few days. Warm salt water rinses (a teaspoon of salt in a cup of warm water) three times a day will reduce discomfort and swelling. A mild pain medication (one tablet of Paracetamol or Ibuprofen every 3-4 hours) should ease any residual discomfort.

Don't be concerned if your speech is affected for the first few days. You'll quickly adapt and be speaking normally. You may notice increased salivation. This is because your brain is responding to the new size and shape of your teeth. This should subside to normal in about a week.

Daily brushing and flossing are a must for your new dental work. Daily plaque removal is critical for the long-term success of your new teeth, as are regular cleaning appointments.

Any food or habits that can crack, chip or damage a natural tooth can do the same to your new teeth. Avoid hard foods and substances such as nuts, ice, fingernails, or pencils and sticky toffee. Smoking will stain your new teeth. Minimize or avoid foods that stain such as coffee, red wine, tea and berries.

If you engage in sports let us know so we can make a custom mouthguard. If you grind or clench your teeth, wear the night guard, when sleeping, that we have provided for you. Adjusting to the look and feel of your new smile will take time. If you have any problems or concerns, please call our surgery. We always welcome your questions.